Common Name			
	Scientific Name	Recommended Survey Window*	Consulting Resource Agencies
		A with Assumption	
	Balaenoptera musculus	April - August	NMFS
	Balaenoptera physalus	April - August	NMFS
umpback whale (E)** orth Atlantic right whale (E)**	Megaptera novaeangliae Eubalaena glacialis	April - August April - August	NMFS
			NMFS NMFS
	Balaenoptera borealis Physeter macrocephalus	April - August April - August	
RACHNIDS	Physeler macrocephalus		NMFS
pruce-fir moss spider (E)**	Microhexura montivaga	May - August	USFWS
IRDS	inicione and included		
		Year round; November - March (optimal to observe birds and nest); February -	
ald eagle (BGPA)	Haliaeetus leucocephalus	May (optimal to observe active nesting)	USFWS
iping plover (T&E)	Charadrius melodus	Year round	USFWS
ed-cockaded woodpecker (E)	Picoides borealis	Year round; November - early March (optimal)	USFWS
oseate tern (E)	Sterna dougallii	June - August	USFWS
ufa red knot (T)	Calidris canutus rufa	Year round	USFWS
/ood stork (T)	Mycteria americana	April 15 - July 15	USFWS
ISH/AMPHIBIAN			
tlantic sturgeon (E)**	Acipenser oxyrinchus oxyrinchus	Not required; assume presence in appropriate waters	NMFS
	Notropis mekistocholas	April - June or periods of high flow (tributaries); Year round (large rivers)	USFWS
arolina madtom (E)	Noturus furiosus	May - October	USFWS
iant manta ray $(T)$	Manta birostris	No survey window established at this time, per NOAA Southeast Fisheries Science Center.	NMFS
iant manta ray (T)	Manta birostris Cryptobranchus alleganiensis		
astern Hellbender (PE)	alleganiensis	No survey window established at this time, per USFWS.	USFWS
	Necturus lewisi	November 15 - March 15	USFWS
		No survey window established at this time, per NOAA Southeast Fisheries	
ceanic whitetip shark (T)	Carcharhinus longimanus	Science Center.	NMFS
	Percina rex	Year round; April - June (optimal)	USFWS
hortnose sturgeon (E)**	Acipenser brevirostrum	Not required; assume presence in appropriate waters	NMFS
potfin chub (T)	Cyprinella monacha	September - November (tributaries); Year round (large rivers)	USFWS
/accamaw silverside (T)	Menidia extensa	Year round	USFWS
ISECTS			
	Bombus affinis	May - September	USFWS
aint Francis' satyr (E)	Neonympha mitchellii francisci	May 5 - June 6 and July 26 - August 21	USFWS
AMMALS		Inay 3 - June 6 and July 26 - August 21	031 103
AMMALS		[May - October (trapping); coldest days in coldest winter months (nest box	
arolina northern flying squirrel (E)	Glaucomys sabrinus	surveys)	USFWS
ray bat (E), Indiana bat (E), Northern	-	West: Bridge surveys: May 1- October 1, concrete/metal, culvert surveys	
	Myotis grisescens, Myotis sodalis,	(3'Hx60'L min): hibernation range = Dec 15 - Feb 15, year round range = no	
sk), Tricolored bat (Proposed E) - Species		culvert surveys *** East: Structure surveys: year	
overed in western Programmatic BO	Perimyotis subflavus	round***	USFWS
ed wolf (E)**	Canis rufus	Year round	USFWS
irginia big-eared bat (E)	Corynorhinus townsendii virginianus	Structure Checks: May 15 - August 15***	USFWS
/est Indian manatee (E)**	Trichechus manatus	Year round	USFWS
USSELS			
ppalachian elktoe (E)	Alasmidonta raveneliana	March 1 - November 1 (optimal)	USFWS
tlantic pigtoe (T)	Fusconaia masoni	March 1 - November 1 (optimal)	USFWS
arolina heelsplitter (E)	Lasmigona decorata	March 1 - November 1 (optimal)	USFWS
umberland bean pearlymussel (E)	Villosa trabalis	March 1 - November 1 (optimal)	USFWS
warf wedgemussel (E)	Alasmidonta heterodon	March 1 - November 1 (optimal)	USFWS
ames spinymussel (E)	Pleurobema collina	March 1 - November 1 (optimal)	USFWS
ttlewing pearly mussel (E)	Pegias fabula	March 1 - November 1 (optimal)	USFWS
umberland moccasinshell (PT)	Medionidus conradicus	March 1 - November 1 (optimal)	USFWS
ennessee pigtoe (PE)	Pleuronaia barnesiana	March 1 - November 1 (optimal)	USFWS
ennessee clubshell (PE)	Pleurobema oviforme	March 1 - November 1 (optimal)	USFWS
ongsolid (T)	Fusconaia subrotunda	March 1 - November 1 (optimal)	USFWS
an riffleshell (E)**	Epioblasma florentina walkeri walkeri	March 1 - November 1 (optimal)	USFWS or NCWRC must request surve
reen floater (PT)	Lasmigona subviridus	March 1 - November 1 (optimal)	USFWS
ar River spinymussel (E)	Elliptio steinstansana	March 1 - November 1 (optimal)	USFWS
ellow lance (T)	Elliptio lanceolata	March 1 - November 1 (optimal)	USFWS
EPTILES			
merican alligator T(S/A)**	Alligator mississippiensis	Year round (only warm days in winter)	USFWS
• • • •	Glyptemys muhlenbergii	April 1 - October 1 (visual surveys)	USFWS or NCWRC must request surve
reen sea turtle (T)**	Chelonia mydas	April - August for beach surveys	USFWS and NMFS
awksbill sea turtle (E)**	Eretmochelys imbricata	April - August for beach surveys	USFWS and NMFS
emp's Ridley sea turtle (E)**	Lepidochelys kempii	April - August for beach surveys	USFWS and NMFS
eatherback sea turtle (E)**	Dermochelys coriacea	April - August for beach surveys	USFWS and NMFS
oggerhead sea turtle (T)**	Caretta caretta	April - August for beach surveys	USFWS and NMFS
NAILS			
oonday globe snail (T)	Petera clarki nantahala	April - September (or prior to first fall frost); April - May (optimal)	USFWS
	Planorbella magnifica	No survey window established at this time, per USFWS.	USFWS
- J			1
rotected animal species surveys romain valid for two w		some as the second of a second approximation with the appropriate resource agency.	
Protected animal species surveys remain valid for two ye	· · · · ·		
NCDOT surveys are not regularly performed for these s	species. Species may be listed as T(S/A), or field sur	veys are too difficult or impractical (record searches and/or resource agency contact are used in lieu of f	
NCDOT surveys are not regularly performed for these s	species. Species may be listed as T(S/A), or field sur t on results of bat structure checks or USFWS require	ements. <u>Mist Netting Surveys</u> : <b>Divisions 1-8</b> : Coastal Plain <sup>†</sup> , April 15 - September 15; Piedmont <sup>†</sup> , May	